

Research Article

PART BOUNDARIES ALTER THE PERCEPTION OF TRANSPARENCY

Manish Singh and Donald D. Hoffman

Department of Cognitive Sciences, University of California, Irvine

Abstract—The perception of transparency is a remarkable feat of human vision: A single stimulation at the retina is interpreted as arising from two (or more) distinct surfaces, separated in depth, in the same visual direction. This feat is intriguing because physical transparency is neither necessary nor sufficient for phenomenal transparency. Many conditions for phenomenal transparency have been studied, including luminance, chromaticity, stereo depth, apparent motion, and structure from motion. Figural conditions have also been studied, primarily by Gestalt psychologists, resulting in descriptive laws. Here we extend, and make precise, these laws using the genericity principle and the minima rule for part boundaries. We report experiments that support the psychological plausibility of these refinements. The results suggest that the formation of visual objects and their parts is an early process in human vision that can precede the representation of transparency.

Figure 1a shows two opaque gray rectangles, one on a dark background and the other on a light background. If the two gray rectangles are moved, so that their edges coincide with each other and with the lightness border (as in Fig. 1b), an observer will see not two opaque gray rectangles, as before, but a single large transparent filter, in front of the divided background. This demonstration shows that physical transparency is not necessary for phenomenal transparency. Also, in Figure 2a, an observer does not see transparency, even though this display might be produced by a transparent filter placed over the bicolored background. Thus, physical transparency is not sufficient for phenomenal transparency.

What conditions determine when transparency will be seen? The most extensively studied conditions for phenomenal transparency are those involving achromatic luminance (Beck, Prazdny, & Ivry, 1984; Gerbino, Stultiens, Troost, & de Weert, 1990; Kanizsa, 1979; Metelli, 1974; Metelli, Da Pos, & Cavedon, 1985), and these have also been extended to the chromatic domain (Da Pos, 1989; D'Zmura, Colantoni, Knoblauch, & Laget, 1997). For example, if the two gray rectangles in Figure 1b are interchanged (as in Fig. 1c), or if both gray rectangles are given the same luminance (as in Fig. 1d), then the perception of transparency is lost. It has also been shown that the perception of transparent surfaces interacts with stereo depth (Nakayama & Shimojo, 1992; Nakayama, Shimojo, & Ramachandran, 1990), subjective contours (Cicerone & Hoffman, 1991; Nakayama et al., 1990), apparent motion (Cicerone & Hoffman, 1991; Cicerone, Hoffman, Gowdy, & Kim, 1995; Shipley & Kellman, 1994, 1997), and structure from motion (Kersten, Buelthoff, Schwartz, & Kurtz, 1992).

Address correspondence to Manish Singh, Perceptual Science Group, Department of Brain and Cognitive Sciences, MIT, Cambridge, MA 02138; e-mail: manish@psyche.mit.edu.




Fig. 1. Luminance conditions for transparency. In (a), the two small gray rectangles are not connected and appear opaque. In (b), these rectangles are placed next to each other and look like a single transparent filter. In (c), the two rectangles are reversed and no longer look transparent. In (d), they are the same shade of gray and again do not look transparent.

It is clear that in displays like those in Figure 1, apart from luminance conditions, certain properties of shape must also be satisfied in order for transparency to be seen. For example, in Figures 1a, 2a, and 2b, transparency is not seen. These displays violate Kanizsa's (1979) *topological condition*: The two gray regions that are to form the transparent surface must be in contact with each other, and each must make contact with only one of the two background regions. A figural condition suggested by Kanizsa (1979) and Metelli (1974) is *discontinuity of direction*. Examples (from Kanizsa, 1979, pp. 158–161) of discontinuity of direction are shown in Figures 2c and 2d. As these displays indicate, by discontinuity of direction, Kanizsa meant two things: discontinuity in the direction of the contour of the filter (as in Fig. 2c) and discontinuity in the direction of the line dividing the background (as in Fig. 2d). Kanizsa gave Figure 2c as an example in which transparency is blocked, and Figure 2d as an example in which transparency is not blocked, by the discontinuity of direction. In this article, we consider only the case of discontinuity of the filter. The experiments we report here suggest two explanations, based on more recent work in vision, that can be cast in precise mathematical terms and that refine the discontinuity explanation. The first is more general, and is based on the




Fig. 2. Kanizsa and Metelli's figural conditions for transparency. In (a) and (b), Kanizsa's (1979) topological condition is violated, and no transparency is seen. In (c), discontinuities of direction destroy transparency, but in (d), they do not.

principle of genericity (e.g., Binford, 1981). The second is based on the *minima rule* (Hoffman & Richards, 1984) for parsing visual shapes and on a part-saliency rule that builds on the minima rule (Hoffman & Singh, 1997). These two explanations are not mutually exclusive, but complement each other.

THE GENERICITY PRINCIPLE

Human vision's interpretations about the visual environment are typically underconstrained by the information available at the retinal images: Countless interpretations are always consistent with any given image or set of images. To deal with this problem, human vision uses various constraints on possible interpretations, and is thus able to reach unique or nearly unique interpretations. The principle of genericity provides one powerful such constraint. In its simplest form, this principle says to reject unstable interpretations of visual stimuli. An unstable interpretation is one that, if perturbed slightly (e.g., with regard to its position relative to the observer), would lead to a qualitative change (e.g., a change in the topological or first-order differential structure) in the projected image.

For example, consider Figure 3a, which shows a Necker cube. Observers readily perceive this illustration as a cube in three dimensions. In Figure 3b, however, the perception of a cube is lost; the figure looks more like a flat pinwheel. In fact, this image is also the projection of a cube, albeit from a special viewing position—one in which two opposite vertices of the cube are perfectly aligned. This viewing position is nongeneric, however, because even a slight change in the viewing position would change the topological structure of the image: For example, the image in Figure 3c has seven connected regions, whereas the image in Figure 3b has six. Because interpreting




Fig. 3. Demonstration of the principle of genericity. In (a), an observer sees a cube in three dimensions. In (b), an observer sees a flat pinwheel and not a cube. The reason, according to the principle of genericity, is that the view in (b) is nongeneric: A slight change in viewpoint would alter the topological structure of the image, as in (c).

Figure 3b as a cube requires assuming a nongeneric (i.e., unstable) viewing position, human vision rejects this interpretation, and observers therefore see Figure 3b as flat.¹

The principle of genericity has been applied, successfully, to provide theories of various visual capacities, including the three-dimensional (3-D) interpretation of line drawings (Binford, 1981; Lowe & Binford, 1985), the perception of subjective contours (Albert & Hoffman, 1995, in press), the perception of object parts (Biederman, 1987; Hoffman & Richards, 1984; Hoffman & Singh, 1997), the perception of shape from shading (Freeman, 1994), and the phenomenon of color constancy (Brainard & Freeman, 1997). It has also been incorporated into formal Bayesian models of visual perception (Freeman, 1996).

To see the role of genericity in the perception of transparency, consider the display in Figure 4a, in which an observer perceives a circular transparent filter over a bicolored background. Figure 4b introduces two concave cusps that fall precisely on the lightness border, and the perception of transparency is greatly reduced. According to the genericity explanation, if there were a transparent filter in front of the divided background, it would take a special viewing position to make the extrema of curvature on the filter align precisely with the lightness border; hence, the interpretation of the transparent filter is nongeneric. Therefore, the luminance change should be interpreted as a reflectance change (i.e., due not to transparency but to different surfaces).

THE MINIMA RULE

There is now growing evidence that human vision represents the shapes of objects in terms of component parts, and the spatial relationships between these parts (Baylis & Driver, 1995a, 1995b; Biederman, 1987; Braunstein, Hoffman, & Saidpour, 1989; Hoffman & Richards, 1984; Hoffman & Singh, 1997; Marr & Nishihara, 1978; Singh, Seyranian, & Hoffman, in press). From a computational perspective, part-based representations of shape provide an efficient way to deal with occluded objects and with articulated objects that do not have fixed shapes—both of which are problems for traditional theories of object recognition, such as template theories and Fourier models. Indeed, recent experimental evidence suggests not only that human

1. One might argue that it is the symmetry of the pinwheel interpretation that is responsible for the perceived flatness. But one gets the same effect with nonregular solids—for which the flat interpretation is not symmetric. Hence, symmetry fails to provide a general explanation of the phenomenon (see Albert & Hoffman, 1995; Kanizsa, 1979, pp. 105–106).




Fig. 4. The role of genericity and the minima rule in the perception of transparency. In (a), an observer sees a circular transparent filter overlying a bicolored background. In (b), two concave cusps are precisely aligned with the luminance border, and these greatly reduce the perception of transparency.

vision parses shapes into parts, but also that it does so quickly, perhaps preattentively (Baylis & Driver, 1995a, 1995b; Driver & Baylis, 1995; Hoffman & Singh, 1997).

Hoffman and Richards (1984) have argued that human vision uses general computational rules to parse objects into parts. Their minima rule defines part boundaries. Because it is expressed solely in the language of differential geometry, it applies quite generally. For a silhouette, the minima rule gives negative minima of curvature as boundary points on the contour of the silhouette. For a 3-D object, it gives loci of negative minima of the principal curvatures as boundary curves on the surface of the object. The *cosine surface* in Figure 5 nicely demonstrates the minima rule. In this illustration, an observer sees circular hills separated by valleys. The boundaries between one hill-shaped part and the next are marked by dashed contours; these are the negative minima of the principal curvatures. If you turn the figure upside down, the figure-ground reversal changes the negative minima to positive maxima, and vice versa. This causes the part boundaries to shift to the new negative minima, and so you now see new parts. The dashed contours that before sat between hills now sit on top of hills.

We (Hoffman & Singh, 1997) have proposed a part-salience rule: The sharper the negative minima of curvature, the more salient the part boundaries. Consider, for example, the Schröder staircase in Figure 6a. This can be seen either as a normal, ascending staircase (the *upright* interpretation) or as a strange, inverted staircase (the *upside-down* interpretation). Observers usually prefer the *upright* interpretation. But in Figure 6b, the staircase has been altered so that the *upside-down* interpretation has more salient part boundaries than the *upright* interpretation. For the *upright* interpretation, the negative minima of curvature (and hence the part boundaries) are the smooth bends along the staircase. If the observer switches figure and ground to see the other (upside-down) interpretation, concave and convex reverse, and the new negative minima are along the sharp bends along the staircase. Hence, the part boundaries for the *upside-down* interpretation are more salient, and observers tend to see the upside-down staircase in Figure 6b, despite the usual preference to see figure below rather than overhead.²

2. This argument is based on our (Hoffman & Singh, 1997) *hypothesis of salient figures*: Other things being equal, human vision prefers that assignment of figure and ground which leads to the figure side having the more salient parts.




Fig. 5. The cosine surface. The parts you see on this surface depend on the orientation in which you view it.




Fig. 6. Demonstration of the part-salience rule. The Schröder staircase in (a) can be seen either as an upright, ascending staircase or as a strange, inverted staircase. Usually there is a preference to see the upright interpretation. But in (b), the staircase is modified so that the inverted interpretation has the more salient part boundaries. As a result, an observer is more likely to see the inverted interpretation in this case.

To see the role of part boundaries in the perception of transparency, consider again the display in Figure 4b. According to the minima-rule explanation, the sharp negative minima of curvature on the filter indicate two distinct parts. Hence, the change in luminance at the part boundaries is interpreted not as transparency, but as different parts of an object having different reflectances.

The difference between the part-boundary and the genericity explanations is that the part-boundary rule predicts a difference between positive maxima and negative minima of curvature (i.e., points of locally greatest magnitude of curvature in convex and concave regions, respectively), whereas the genericity principle does not. Specifically, the minima rule predicts that the presence of negative minima should impair the perception of transparency more than the presence of positive maxima of comparable strength.

EXPERIMENTS

We ran two experiments to demonstrate the genericity effect and the minima-rule effect in the perception of transparency. The first experiment pit filters with no extrema (see Fig. 4a) against those with extrema (see Fig. 7) aligned with the lightness border to explore the genericity effect; and it pit negative minima against positive maxima to explore the minima-rule effect. In addition, it looked at the effect of smoothing and turning angle at the extrema—in light of our (Hoffman & Singh, 1997) theory of part salience. In the second experiment, we sought to provide further support for the genericity explanation by looking at the effect of displacing the curvature extrema of the filters from the lightness border.

Experiment 1

Method

Subjects. The subjects were 8 graduate students at the University of California, Irvine. They were naive to the purposes of the experiment.




Fig. 7. Eight of the stimuli used in Experiment 1. The filters in (a) through (d) have negative minima of curvature, perfectly aligned with the lightness border, and the filters in (e) through (h) have positive maxima. The extrema in (a), (b), (e), and (f) are cusps, and the extrema in (c), (d), (g), and (h) have the high level of smoothing. The filters in (a), (c), (e), and (g) have the lowest magnitude of turning angle (Level 1, 42°), and the filters in (b), (d), (f), and (h) have the highest magnitude of turning angle (Level 4, 132°).

Stimuli. The stimuli were 25 displays like the ones in Figure 7. The CIE (Commission Internationale de l'Eclairage) coordinates and luminance values of the four regions were as follows: lightest gray— $x = 0.273$, $y = 0.269$, luminance = 46.2 cd/m²; light gray— $x = 0.268$, $y = 0.264$, luminance = 23.0 cd/m²; dark gray— $x = 0.248$, $y = 0.234$, luminance = 3.59 cd/m²; black—luminance = 0 cd/m². The displays were viewed at a distance of 0.5 m, and each was about 15° tall and 20° wide. One display had a circle as the transparent filter (see Fig. 4a). Twelve displays had negative minima of curvature that were perfectly aligned with the lightness border (as in Figs. 7a–7d). Twelve displays had positive maxima of curvature, perfectly aligned with the lightness border (as in Figs. 7e–7h).

A wiggle was drawn down the middle of the lightness border to suppress the perception of a crease in 3-D, which is striking in the stimuli with strong negative minima and positive maxima. In a pilot study, subjects reported that this 3-D crease interfered with their judgments of transparency. The length of the wiggle was adjusted in each display so that it stopped at about 1° of visual angle from the X- and ψ -junctions.

Design. There were three independent variables: sign of curvature at the extrema, turning angle at the extrema, and level of smoothing at the extrema. All factors were run within subjects. The sign of curvature had two levels: positive maxima of curvature and negative minima of curvature. The turning angle at the extrema had four levels, labeled 1, 2, 3, and 4 (defined later). The smoothing had three levels: cusp, low level of smoothing, and high level of smoothing. (Fig. 7 shows eight of the stimuli used.) All stimuli were part of this 2 × 4 × 3 factorial design, except for the stimulus with the circle—for which smoothing is inapplicable. Hence, there was a total of 25 (24 + 1) stimuli. Each stimulus was presented 10 times, resulting in a total of 250 experimental trials per subject. These were preceded by 50 practice trials. Whether the more luminous side of the bicolored background appeared on the left or on the right of each display was counterbalanced.

For the stimuli with cusps, the four levels of turning angle were 42°, 72°, 102°, and 132°. Their smoothed versions were created by taking the cusp version and replacing a region of the contour around the cusp with an arc from the tip of an ellipse. The dimensions of this ellipse were 1.5° × 1.1° of visual angle in the low-smoothing case and 3.1° × 2.3° of visual angle in the high-smoothing case.

Apparatus. The figures were displayed on a monitor measuring 1,024 by 768 pixels by a Macintosh Quadra computer using the SuperLab program. Subjects used a keyboard to respond.

Procedure. Subjects were instructed that on each trial, they would see a figure composed of four regions with different shades of gray. They were to judge whether the two regions in the center were transparent, using a scale from 1 to 7 (1 = “no transparency; I see four opaque regions”; 4 = “moderate transparency”; 7 = “strong transparency; I see a gray filter over a black and white background”).

The displays were presented in random order. Each trial consisted of a fixation dot for 500 ms, a transparency display for 2 s, and then a response screen that asked the subject to rate the transparency of the display from 1 to 7. This screen remained until the subject responded.

Results and discussion

Figure 8 shows the results of the first experiment. Transparency ratings were significantly lower for both the negative-minima and positive-maxima cases, as compared with the circle case, $F(1, 7) = 177.32, p < .0001$. This result supports the genericity explanation, because a transparent interpretation of a display in which extrema (positive maxima or negative minima) of curvature align with the lightness border would be nongeneric.

The results also support the minima-rule explanation, because the transparency ratings were significantly lower for negative minima than for positive maxima, $F(1, 7) = 30.048, p < .001$. As mentioned earlier, this result is not predicted by the genericity explanation. Furthermore, there was a main effect of the smoothing level, $F(2, 14) = 15.353, p < .0005$: Transparency ratings were lower for cusp extrema than for smooth extrema. This result supports our (Hoffman & Singh, 1997) theory of part salience, according to which cusp boundaries are more salient than smooth ones. There was also a main effect of turning angle, $F(3, 21) = 9.635, p < .0005$, which is another factor in the theory of part salience (i.e., larger turning angles are indicative of more salient part boundaries).

In sum, these results show that the “discontinuities” explanation of Kanizsa and Metelli can be extended in two ways:

1. Neither tangent discontinuities nor discontinuities of higher derivatives are necessary to block the perception of transparency because the ratings of transparency go down even when the extrema of curvature are smooth. In other words, in order to have a loss of phenomenal transparency, it is sufficient to have strong negative minima or positive maxima of curvature that align with the lightness border, even if these extrema have continuous higher order derivatives.
2. The discontinuities explanation does not predict a difference between negative minima and positive maxima of curvature. The minima rule does predict this difference.

Experiment 2

The purpose of the second experiment was to further support the genericity explanation, by showing that it is indeed the precise alignment of the extrema of curvature with the lightness border that leads to the decline in transparency ratings. We predicted that if the extrema were displaced from the lightness border, then transparency ratings would increase.




Fig. 8. Results of Experiment 1. Mean transparency ratings are shown as a function of sign of curvature (negative minima, positive maxima), turning angle (Levels 1, 2, 3, 4), and degree of smoothing (cusp, low smoothing, high smoothing). The 0 corresponds to the display with the circular filter (see Fig. 4a). Transparency ratings were made on a scale from 1 (*no transparency*) to 7 (*strong transparency*).

Method

Subjects. The subjects were 8 graduate students at the University of California, Irvine. They were naive to the purposes of the experiment.

Stimuli. The stimuli were 18 transparency-type displays that had the same respective luminance values as the displays in Experiment 1. The displays were viewed at a distance of 0.5 m, and each was about 15° tall and 20° wide. Six of the displays were taken from Experiment 1, namely, the 6 displays with the most extreme turning angles (both positive and negative). These 6 displays were then modified by displacing, by two different amounts, the extrema of curvature from the lightness border. The *small* displacement was a displacement of 0.25° (see Figs. 9a, 9c, 9e, and 9g), and the *large* displacement was a displacement of 2° (see Figs. 9b, 9d, 9f, and 9h).

Design. There were three independent variables: sign of curvature at the extrema, level of smoothing at the extrema, and the displacement




Fig. 9. Eight of the stimuli used in Experiment 2. The filters in (a) through (d) have negative minima of curvature on their outlines, and the filters in (e) through (h) have positive maxima. The extrema in (a), (b), (e), and (f) are cusps, and the extrema in (c), (d), (g), and (h) have the high level of smoothing. In (a), (c), (e), and (g), the extrema on the filters have the small level of displacement (0.25°) from the lightness border, and in (b), (d), (f), and (h), the extrema have the large level of displacement (2°).

of the extrema from the lightness border. The sign of curvature had two levels: positive maxima of curvature and negative minima of curvature. The smoothing had three levels: cusp, low level of smoothing, and high level of smoothing. And the displacement had three levels: zero displacement, small displacement, and large displacement. The variables formed a $2 \times 3 \times 3$ factorial design. All variables were run within subjects. Each display was presented 10 times, resulting in a total of 180 experimental trials per subject. These were preceded by 36 practice trials.

The following variables were counterbalanced: whether the more luminous side of the bicolored background appeared on the left or on the right of the display and whether the extrema were displaced to the left or to the right of the line dividing the bicolored background.

Apparatus. The apparatus was the same as in Experiment 1.

Procedure. The instructions were precisely the same as in Experiment 1. The displays were presented in random order. Each trial was structured the same way as in Experiment 1.

Results and discussion

Figure 10 shows the results for the second experiment. As predicted, there was a main effect of the level of displacement, $F(2, 14) = 34.946, p < .0001$. In fact, for displays with large displacements, mean ratings came back up almost as high as the highest ratings in Experiment 1 (i.e., for the display with the circle). There was also a main effect of smoothing, $F(2, 14) = 9.194, p < .003$, but no main effect of the sign of curvature, $F(1, 7) = 0.131, p > .7$.

There was a significant interaction between smoothing and sign of curvature, $F(2, 14) = 4.774, p < .03$. Post hoc analysis revealed that for displays with smooth extrema, transparency ratings were not significantly different between positive maxima and negative minima. However, for displays with cusps, transparency ratings were significantly higher for positive maxima than for negative minima.

There was also an interaction between smoothing and displacement of extrema, $F(4, 28) = 17.956, p < .0001$. Post hoc analysis revealed that for large displacements, there was no significant effect of smoothing level, but for zero and small displacements, the ratings for the cusp displays were significantly lower than the ratings for the smoothed displays.

These results confirm the predictions of the genericity principle: It is indeed the precise alignment of the extrema of curvature with the lightness border that is responsible for the decline in transparency ratings when the displacement is zero.

AN ALTERNATIVE HYPOTHESIS

An alternative angle hypothesis might be advanced to explain our results. The role of X- and ψ -junctions is well-known in the transparency literature (see, e.g., Anderson, 1997; Kersten, 1991). It might be argued that the strength of perceived transparency depends not on the genericity principle and the minima rule, but rather on the angle between the filter contour and the lightness border (dividing the bicolored background) at each X- or ψ -junction. For example, in Figure 4a, the filter contour is orthogonal to the lightness border, whereas in Figure 4b, the filter contour meets the lightness border at a sharp angle. It is perhaps this difference in angle that is responsible for the results obtained in Experiments 1 and 2.




Fig. 10. Results of Experiment 2. Mean transparency ratings are shown as a function of sign of curvature (negative minima, positive maxima), degree of smoothing (cusp, low smoothing, high smoothing), and level of displacement of the extrema from the lightness border (none, small, large). Transparency ratings were made on a scale from 1 (*no transparency*) to 7 (*strong transparency*).

Recall, however, that the results of Experiment 1 showed a significant difference between displays with negative minima and positive maxima of curvature, even though the angle magnitudes were controlled. For example, displays with negative-minima cusps were consistently rated lower in transparency than the corresponding displays with positive-maxima cusps, even though the angles between the contour and the lightness border were the same in both cases (see Fig. 8).

Therefore, an explanation based on the angle hypothesis is insufficient to explain our results.

However, to test the angle hypothesis directly, we ran a control experiment using the three displays shown in Figure 11: One had a horizontally oriented ellipse (so that its contour was orthogonal to the lightness border); another had an ellipse oriented at an angle (so its contour made an angle of 30° with the lightness border); and the third




Fig. 11. The three stimuli used in the control experiment. The stimuli were designed to separate the genericity and minima-rule hypotheses from the angle hypothesis. The genericity and minima-rule hypotheses predict that transparency ratings should be high for the stimuli in the top two panels, and low for the stimulus in the bottom panel. The angle hypothesis predicts that transparency ratings should be high only for the stimulus in the top panel, and should be low for the stimuli in the bottom two panels.

was a version of the negative-minima cusp stimuli used in Experiments 1 and 2—with the constraint that the contour made an angle of 30° with the lightness border. All luminance values were the same as in the first two experiments. We counterbalanced two variables: whether the darker side appeared to the left or to the right and whether the slope of the oblique ellipse was positive or negative.

Five subjects performed the same transparency-rating task as in the first two experiments. We found a significant effect of display type, $F(2, 8) = 26.74, p < .001$. Subjects consistently gave high ratings to the

displays with the horizontal ellipse ($M = 5.15, SE = 0.421$) and the oblique ellipse ($M = 6.23, SE = 0.384$), despite the difference in angles, and they consistently gave low ratings to the display in which the negative minima were aligned with the lightness border ($M = 1.83, SE = 0.318$), even though the angles for this display were the same as those for the display with the oblique ellipse. Post hoc analysis revealed that the mean transparency ratings for the displays with the horizontal ellipse and the oblique ellipse did not differ significantly from each other, but did differ significantly from the mean rating for the display with the negative minima. These results disconfirm the angle hypothesis and support the genericity and minima-rule hypotheses.

CONCLUSIONS

We have proposed that the genericity principle, the minima rule, and a part-salience rule provide a rigorous refinement of the Gestalt figural conditions for the perception of transparency. The experiments reported here support the psychological plausibility of these refinements.

The experiments also support the idea that human vision constructs various properties of visual objects in a highly coordinated fashion (Hoffman, 1998; Singh & Hoffman, 1997). When the central regions in transparency-type displays are seen as a one-part object, they are perceived as being transparent and having uniform reflectance, but when they are seen as two parts of an object, they are perceived as being opaque, with the two parts having different reflectances.

The experimental results suggest that the formation of visual objects and their parts can precede the representation of transparency. This may be surprising because transparency seems to be such a basic visual property. However, there is now psychophysical evidence suggesting that part formation is an early visual process (Hoffman & Singh, 1997), and possibly preattentive (Baylis & Driver, 1995a, 1995b; Driver & Baylis, 1995). So it is perhaps not surprising to find that other visual properties, such as transparency, depend on it.

Acknowledgments—For discussions and suggestions, we thank Bill Batchelder, Bruce Bennett, Myron Braunstein, Mike D’Zmura, and Grace Mack. And we thank two anonymous reviewers for helpful comments on previous versions of this article.

REFERENCES

- Albert, M.K., & Hoffman, D.D. (1995). Genericity in spatial vision. In R.D. Luce, M. D’Zmura, D.D. Hoffman, G.J. Iverson, & A.K. Romney (Eds.), *Geometric representations of perceptual phenomena* (pp. 95–112). Mahwah, NJ: Erlbaum.
- Albert, M.K., & Hoffman, D.D. (in press). Generic visions: General position assumptions in visual perception. *Scientific American*.
- Anderson, B.L. (1997). A theory of illusory lightness and transparency in monocular and binocular images: The role of contour junctions. *Perception*, *26*, 419–453.
- Baylis, G.C., & Driver, J. (1995a). Obligatory edge assignment in vision: The role of figure and part segmentation in symmetry detection. *Journal of Experimental Psychology: Human Perception and Performance*, *21*, 1323–1342.
- Baylis, G.C., & Driver, J. (1995b). One-sided edge assignment in vision: 1. Figure-ground segmentation and attention to objects. *Current Directions in Psychological Science*, *4*, 140–146.
- Beck, J., Pradzny, K., & Ivry, R. (1984). The perception of transparency with achromatic colors. *Perception & Psychophysics*, *35*, 407–422.
- Biederman, I. (1987). Recognition-by-components: A theory of human image understanding. *Psychological Review*, *94*, 115–147.

Part Boundaries and Transparency

- Binford, T.O. (1981). Inferring surfaces from images. *Artificial Intelligence*, *17*, 205–244.
- Brainard, D.H., & Freeman, W.T. (1997). Bayesian color constancy. *Journal of the Optical Society of America A*, *14*, 1393–1411.
- Braunstein, M.L., Hoffman, D.D., & Saidpour, A. (1989). Parts of visual objects: An experimental test of the minima rule. *Perception*, *18*, 817–826.
- Cicerone, C.M., & Hoffman, D.D. (1991). *Dynamic neon colors: Perceptual evidence for parallel visual pathways* (UCI Mathematical Behavioral Sciences Memo No. 91–22). Irvine, CA: Institute of Mathematical Behavioral Sciences.
- Cicerone, C.M., Hoffman, D.D., Gowdy, P.D., & Kim, J.S. (1995). The perception of color from motion. *Perception & Psychophysics*, *57*, 761–777.
- Da Pos, O. (1989). *Trasparenze*. Padua, Italy: Iacone.
- Driver, J., & Baylis, G.C. (1995). One-sided edge assignment in vision: 2. Part decomposition, shape description, and attention to objects. *Current Directions in Psychological Science*, *4*, 201–206.
- D’Zmura, M., Colantoni, P., Knoblauch, K., & Laget, B. (1997). Color transparency. *Perception*, *26*, 471–492.
- Freeman, W.T. (1994). The generic viewpoint assumption in a framework for visual perception. *Nature*, *368*, 542–545.
- Freeman, W.T. (1996). The generic viewpoint assumption in a Bayesian framework. In D.C. Knill & W.A. Richards (Eds.), *Perception as Bayesian inference* (pp. 365–389). New York: Cambridge University Press.
- Gerbino, W., Stulmans, C.I., Troost, J.M., & de Weert, C.M. (1990). Transparent layer constancy. *Journal of Experimental Psychology: Human Perception and Performance*, *16*, 3–20.
- Hoffman, D.D. (1998). *Visual intelligence: How we create what we see*. New York: Norton.
- Hoffman, D.D., & Richards, W.A. (1984). Parts of recognition. *Cognition*, *18*, 65–96.
- Hoffman, D.D., & Singh, M. (1997). Saliency of visual parts. *Cognition*, *63*, 29–78.
- Kanizsa, G. (1979). *Organization in vision*. New York: Praeger.
- Kersten, D. (1991). Transparency and the cooperative computation of scene attributes. In M.S. Landy & J.A. Movshon (Eds.), *Computational models of visual processing* (pp. 209–228). Cambridge, MA: MIT Press.
- Kersten, D., Buelhoff, H.H., Schwartz, B.L., & Kurtz, K.J. (1992). Interaction between transparency and structure from motion. *Neural Computation*, *4*, 573–589.
- Lowe, D.G., & Binford, T.O. (1985). The recovery of three-dimensional structure from image curves. *IEEE Transactions on PAMI*, *7*, 320–326.
- Marr, D., & Nishihara, K. (1978). Representation and recognition of the spatial organization of three-dimensional shapes. *Proceedings of the Royal Society of London B*, *200*, 269–294.
- Metelli, F. (1974). The perception of transparency. *Scientific American*, *230*, 90–98.
- Metelli, F., Da Pos, O., & Cavedon, A. (1985). Balanced and unbalanced, complete and partial transparency. *Perception & Psychophysics*, *38*, 354–366.
- Nakayama, K., & Shimojo, S. (1992). Experiencing and perceiving visual surfaces. *Science*, *257*, 1357–1363.
- Nakayama, K., Shimojo, S., & Ramachandran, V.S. (1990). Transparency: Relation to depth, subjective contours, luminance, and neon color spreading. *Perception*, *19*, 497–513.
- Shipley, T.F., & Kellman, P.J. (1994). Spatio-temporal boundary formation—Boundary, form, and motion perception from transformations of surface elements. *Journal of Experimental Psychology: General*, *123*, 3–20.
- Shipley, T.F., & Kellman, P.J. (1997). Spatio-temporal boundary formation: The role of local motion signals in boundary perception. *Vision Research*, *37*, 1281–1293.
- Singh, M., & Hoffman, D.D. (1997). Constructing and representing visual objects. *Trends in Cognitive Sciences*, *1*, 98–102.
- Singh, M., Seyranian, G., & Hoffman, D.D. (in press). Parsing silhouettes: The short-cut rule. *Perception & Psychophysics*.

(RECEIVED 8/20/97; ACCEPTED 2/19/98)